

HYDROCORTISONE SODIUM SUCCINATE for Injection, USP
For Intravenous or Intramuscular Administration

DESCRIPTION


Hydrocortisone sodium succinate for injection, USP is an anti-inflammatory glucocorticoid that contains hydrocortisone sodium succinate, USP as the active ingredient. Hydrocortisone sodium succinate for injection, USP is available in below mentioned package for intravenous or intramuscular administration.

100 mg Plain—Vials containing hydrocortisone sodium succinate, USP equivalent to 100 mg hydrocortisone sodium succinate, USP. The vial contains 100 mg of hydrocortisone sodium succinate, USP in 10 mL of sterile, nonpyrogenic, isotonic, clear, colorless, and odorless solution. The pH of the container diluent (see **DOSEAGE AND ADMINISTRATION: Preparation of Solutions**).

When necessary, the pH was adjusted with sodium hydroxide so that the pH of the reconstituted solution is within the USP specified range of 7 to 8.

The chemical name for hydrocortisone sodium succinate is pregn-4-ene-3,20-dione-21-(3'-carboxy-1'-oxopropyl)-11,17-dihydro-, monosodium salt, (11β)- and its molecular weight is 484.52.

The structural formula is represented below:



Hydrocortisone sodium succinate is a white or nearly white, odorless, hygroscopic amorphous solid. It is very soluble in water and in alcohol, very slightly soluble in acetone, and insoluble in chloroform.

CLINICAL PHARMACOLOGY

Naturally occurring glucocorticoids, hydrocortisone and cortisone, which also have salt-retaining properties, are used as replacement therapy in adrenocortical deficiency states. Their synthetic analogs are primarily used for their anti-inflammatory effects in disorders of many organ systems.

Hydrocortisone sodium succinate has the same metabolic and anti-inflammatory actions as hydrocortisone. When given parenterally in equimolar quantities, the two compounds are equally effective. The primary effect of hydrocortisone sodium succinate is the inhibition of hydrocortisone permits the intravenous, intramuscular, or intracortical administration of high doses of hydrocortisone in a small volume of diluent and is particularly useful where high blood levels of hydrocortisone are required rapidly. Following the intravenous injection of hydrocortisone sodium succinate, demonstrable effects are evident within one hour and persist for a variable period of time. In patients with adrenocortical deficiency, the blood glucose level of a patient with consistently high blood levels are noted. Injections should be made every 4 to 6 hours. This pattern is also rapidly absorbed when administered intramuscularly and is excreted in a pattern similar to that observed after intravenous injection.

Glucocorticoids cause profound and varied metabolic effects. In addition, they modify the body's immune responses to diverse stimuli.

INDICATIONS AND USAGE

When oral therapy is not feasible, and the strength, dosage form, and route of administration of the drug are important factors in the management of the disease, the use of hydrocortisone sodium succinate for injection is indicated. The following are the **INDICATIONS AND USAGE** of hydrocortisone sodium succinate for injection as indicated in **Labeling** or **Intracortical use** of hydrocortisone sodium succinate for injection as indicated in **Labeling**.

Allergic states: Control of severe or incapacitating allergic conditions intractable to adequate trials of conventional treatment in asthma, atopic dermatitis, contact dermatitis, drug hypersensitivity reactions, serum sickness, transfusion reactions.

Dermatologic diseases: Bullous dermatitis herpetiformis, exfoliative erythroderma, mycosis fungoides, pemphigus, severe erythema multiforme (Stevens-Johnson syndrome).

Endocrine disorders: Primary or secondary adrenocortical insufficiency (hydrocortisone or cortisone is the drug of choice; synthetic analogs may be used in conjunction with mineralocorticoids where applicable in infancy, mineralocorticoid supplementation is of value in congenital adrenal hyperplasia, hyperadrenia associated with cancer, postoperative thyroiditis).

Contraindications and Precautions: Drug interactions, Amphotericin B injection and potassium-depleting agents).

Endocrine: Hypothalamic-pituitary adrenal (HPA) axis suppression; Cushing's syndrome, and

Hydrocortisone sodium succinate for injection is contraindicated for intrathecal administration. Reports of severe medical events have been associated with this route of administration.

Warnings: Series Neurologic Adverse Reactions with Epidural Administration

Reports of severe medical events have been associated with this route of administration. The following are the **WARNINGS** of hydrocortisone sodium succinate for injection as indicated in **Labeling**.

Contraindications: Hydrocortisone sodium succinate for injection is contraindicated in patients with known hypersensitivity to the product and its constituents.

Contraindications and Precautions: Drug interactions, Amphotericin B injection and potassium-depleting agents).

Endocrine: Hypothalamic-pituitary adrenal (HPA) axis suppression; Cushing's syndrome, and

Hydrocortisone sodium succinate for injection is contraindicated for intrathecal administration. Reports of severe medical events have been associated with this route of administration.

Neurologic disorders: Acquired (autoimmune) hemolytic anemia, congenital (erythroid) hypoplastic anemia (Diamond Blackfan anemia), idiopathic thrombocytopenic purpura in adults (intravenous administration may be contraindicated), pure red cell aplasia, select cases of secondary thrombocytopenia.

Miscellaneous: Trichinosis with neurologic or myocardial involvement, tuberculous meningitis and other central nervous system infections, including those with concurrent use of appropriate antimicrobial chemotherapy.

Neurologic disorders: For the palliative management of leukemias and lymphomas.

Nervous System: Cerebral edema associated with primary or metastatic brain tumor, or craniospinal irradiation.

Ophthalmic diseases: Sympathetic ophthalmia, exudative and ocular inflammatory conditions unresponsive to topical corticosteroids.

Respiratory diseases: Berylliosis, fulminating or disseminated pulmonary tuberculosis when used concurrently with appropriate antituberculous chemotherapy, idiopathic eosinophilic pneumonias, symptomatic sarcoidosis.

Rheumatic disorders: As adjunctive therapy for short-term administration (to tide the patient over an acute episode or exacerbation) in acute gouty arthritis, acute rheumatic carditis, acute rheumatic fever, acute rheumatoid arthritis, acute rheumatoid arthritis, acute rheumatoid arthritis (radial cases may require low-dose maintenance therapy). For the treatment of dermatomyositis, temporal arteritis, polymyositis, and systemic lupus erythematosus.

CONTRAINDICATIONS

Hydrocortisone sodium succinate for injection is contraindicated in systemic fungal infections and patients with known hypersensitivity to the product and its constituents.

Intramuscular corticosteroid preparations are contraindicated for idiopathic thrombocytopenic purpura.

Hydrocortisone sodium succinate for injection is contraindicated for intrathecal administration. Reports of severe medical events have been associated with this route of administration.

Warnings: Series Neurologic Adverse Reactions with Epidural Administration

Reports of severe medical events have been associated with this route of administration. The following are the **WARNINGS** of hydrocortisone sodium succinate for injection as indicated in **Labeling**.

Contraindications: Hydrocortisone sodium succinate for injection is contraindicated in patients with known hypersensitivity to the product and its constituents.

Contraindications and Precautions: Drug interactions, Amphotericin B injection and potassium-depleting agents).

Endocrine: Hypothalamic-pituitary adrenal (HPA) axis suppression; Cushing's syndrome, and

Hydrocortisone sodium succinate for injection is contraindicated for intrathecal administration. Reports of severe medical events have been associated with this route of administration.

Warnings: Series Neurologic Adverse Reactions with Epidural Administration

Reports of severe medical events have been associated with this route of administration. The following are the **WARNINGS** of hydrocortisone sodium succinate for injection as indicated in **Labeling**.

Contraindications: Hydrocortisone sodium succinate for injection is contraindicated in patients with known hypersensitivity to the product and its constituents.

Contraindications and Precautions: Drug interactions, Amphotericin B injection and potassium-depleting agents).

Endocrine: Hypothalamic-pituitary adrenal (HPA) axis suppression; Cushing's syndrome, and

Hydrocortisone sodium succinate for injection is contraindicated for intrathecal administration. Reports of severe medical events have been associated with this route of administration.

hypertension. Monitor patients for these conditions with chronic use. Corticosteroids can produce reversible HPAAx suppression with the potential for glucocorticosteroid insufficiency after withdrawal of treatment. Drug-induced secondary adrenocortical insufficiency may be severe, especially after prolonged therapy. During or after treatment, the patient may experience prostration or hypotension if he or she receives a sudden reduction in the dose of corticosteroids. If such symptoms occur, the patient should be placed on a gradual reduction of the dose of corticosteroids. If such symptoms occur, the patient should be placed on a gradual reduction of the dose of corticosteroids. If such symptoms occur, the patient should be placed on a gradual reduction of the dose of corticosteroids.

Immunosuppression and increased risk of infection

Corticosteroids, including hydrocortisone sodium succinate, suppress the immune system and increase the risk of infection with any pathogen, including viral, bacterial, fungal, protozoan, or helminthic pathogens. Corticosteroids can:

- Mask some signs of infection
- Increase the risk of dissemination of infections
- Exacerbate existing infections

Corticosteroid-associated infections can be mild but can be severe and at times fatal. The rate of infectious complications increases with increasing corticosteroid dosages.

Monitor for the development of infection and consider hydrocortisone sodium succinate for injection to be discontinued if infection is suspected. Discontinuation of corticosteroids may result in clinical improvement of Kaposi's sarcoma.

Neurologic:

The lowest possible dose of corticosteroid should be used to control the condition under treatment. When reduction in dosage is possible, the reduction should be gradual.

Since complications of treatment with glucocorticoids are dependent on the size of the dose and the duration of treatment, a risk/benefit decision must be made in each individual case as to the dose and duration of treatment and as to whether daily or intermittent therapy should be used.

Contraindications:

Hydrocortisone sodium succinate for injection is contraindicated in patients with known hypersensitivity to the product and its constituents.

Contraindications and Precautions: Drug interactions, Amphotericin B injection and potassium-depleting agents).

Endocrine: Hypothalamic-pituitary adrenal (HPA) axis suppression; Cushing's syndrome, and

Hydrocortisone sodium succinate for injection is contraindicated for intrathecal administration. Reports of severe medical events have been associated with this route of administration.

Warnings: Series Neurologic Adverse Reactions with Epidural Administration

Reports of severe medical events have been associated with this route of administration. The following are the **WARNINGS** of hydrocortisone sodium succinate for injection as indicated in **Labeling**.

Contraindications: Hydrocortisone sodium succinate for injection is contraindicated in patients with known hypersensitivity to the product and its constituents.

Contraindications and Precautions: Drug interactions, Amphotericin B injection and potassium-depleting agents).

Endocrine: Hypothalamic-pituitary adrenal (HPA) axis suppression; Cushing's syndrome, and

Hydrocortisone sodium succinate for injection is contraindicated for intrathecal administration. Reports of severe medical events have been associated with this route of administration.

Warnings: Series Neurologic Adverse Reactions with Epidural Administration

Reports of severe medical events have been associated with this route of administration. The following are the **WARNINGS** of hydrocortisone sodium succinate for injection as indicated in **Labeling**.

Contraindications: Hydrocortisone sodium succinate for injection is contraindicated in patients with known hypersensitivity to the product and its constituents.

Contraindications and Precautions: Drug interactions, Amphotericin B injection and potassium-depleting agents).

Endocrine: Hypothalamic-pituitary adrenal (HPA) axis suppression; Cushing's syndrome, and

Hydrocortisone sodium succinate for injection is contraindicated for intrathecal administration. Reports of severe medical events have been associated with this route of administration.

Vaccination:

Administration of live or live, attenuated vaccines is contraindicated in patients receiving immunosuppressive doses of corticosteroids. Killed or inactivated vaccines may be administered.

Reports of severe medical events have been associated with the intrathecal route of administration (see **ADVERSE REACTIONS: Neurologic/Psychiatric**).

Ophthalmic:

Use of corticosteroids may produce posterior subcapsular cataracts, glaucoma with possible increased intraocular pressure and optic neuropathy which may result in irreversible loss of vision due to backward bulging of the optic disc. The use of oral corticosteroids is not recommended in the treatment of optic neuritis and may lead to an increase in the risk of new episodes. Corticosteroids should be used cautiously in patients with ocular herpes simplex because of corneal perforation. Corticosteroids should not be used in active ocular herpes simplex.

Kaposi's Sarcoma

Kaposi's sarcoma has been reported to occur in patients receiving corticosteroid therapy. Discontinuation of corticosteroids may result in clinical improvement of Kaposi's sarcoma.

Neurologic:

The lowest possible dose of corticosteroid should be used to control the condition under treatment. When reduction in dosage is possible, the reduction should be gradual.

Since complications of treatment with glucocorticoids are dependent on the size of the dose and the duration of treatment, a risk/benefit decision must be made in each individual case as to the dose and duration of treatment and as to whether daily or intermittent therapy should be used.

Contraindications:

Hydrocortisone sodium succinate for injection is contraindicated in patients with known hypersensitivity to the product and its constituents.

Contraindications and Precautions: Drug interactions, Amphotericin B injection and potassium-depleting agents).

Endocrine: Hypothalamic-pituitary adrenal (HPA) axis suppression; Cushing's syndrome, and

Hydrocortisone sodium succinate for injection is contraindicated for intrathecal administration. Reports of severe medical events have been associated with this route of administration.

Warnings: Series Neurologic Adverse Reactions with Epidural Administration

Reports of severe medical events have been associated with this route of administration. The following are the **WARNINGS** of hydrocortisone sodium succinate for injection as indicated in **Labeling**.

Contraindications: Hydrocortisone sodium succinate for injection is contraindicated in patients with known hypersensitivity to the product and its constituents.

Contraindications and Precautions: Drug interactions, Amphotericin B injection and potassium-depleting agents).

Endocrine: Hypothalamic-pituitary adrenal (HPA) axis suppression; Cushing's syndrome, and

Hydrocortisone sodium succinate for injection is contraindicated for intrathecal administration. Reports of severe medical events have been associated with this route of administration.

Warnings: Series Neurologic Adverse Reactions with Epidural Administration

Reports of severe medical events have been associated with this route of administration. The following are the **WARNINGS** of hydrocortisone sodium succinate for injection as indicated in **Labeling**.

Contraindications: Hydrocortisone sodium succinate for injection is contraindicated in patients with known hypersensitivity to the product and its constituents.

Contraindications and Precautions: Drug interactions, Amphotericin B injection and potassium-depleting agents).

Endocrine: Hypothalamic-pituitary adrenal (HPA) axis suppression; Cushing's syndrome, and

Hydrocortisone sodium succinate for injection is contraindicated for intrathecal administration. Reports of severe medical events have been associated with this route of administration.

